



The  
University  
Of  
Sheffield.

MAS315

SCHOOL OF MATHEMATICS AND STATISTICS

Autumn Semester 2016–17

WAVES

2 hours

*Marks will be awarded for your best FOUR answers. The marks awarded to each question or section of question are shown in italics.*

Please leave this exam paper on your desk  
Do not remove it from the hall

Registration number from U-Card (9 digits)  
to be completed by student

--	--	--	--	--	--	--	--	--

Blank

- 1 A string with uniform mass per unit length  $\rho$  is under a tension  $F$ , and undergoes small transverse vibrations. Let the displacement of the point at distance  $x$  along the string at time  $t$  be  $y(x, t)$ .

(i) Show that

$$\frac{\partial^2 y}{\partial x^2} = \frac{1}{c^2} \frac{\partial^2 y}{\partial t^2}$$

to good approximation, where the constant  $c^2 = F/\rho$ .

(5 marks)

(ii) Derive the general solution (*d'Alembert's solution*) of this equation.

(11 marks)

(iii) In a particular case the string is unbounded in both directions, i.e.  $-\infty < x < \infty$ . At  $t = 0$

$$y(x, 0) = 0, \quad \dot{y}(x, 0) = V \cos(2kx),$$

where  $V$  and  $k$  are constants. Find  $y(x, t)$  and all values of  $x$  for which  $y(x, t) = 0$ . Describe the motion very briefly.

(9 marks)

- 2 A uniform finite string of length  $l$  and mass per unit length  $\rho$  occupies the interval  $0 \leq x \leq l$  and undergoes transverse vibrations with displacement  $y(x, t)$ , where  $c^2 y_{xx} = y_{tt}$ , and  $c^2$  is a constant. The tension in the string is  $\rho c^2$ . You are given that

(a)  $y(0, t) = y(l, t) = 0$ ;

(b)  $y(x, 0) = (h/l^2)x(l-x)$  where  $h$  is a constant;

(c)  $\dot{y}(x, 0) = 0$ ;

(d)  $y(x, t) = \sum_{n=1}^{\infty} a_n \sin\left(\frac{n\pi x}{l}\right) \cos\left(\frac{n\pi ct}{l}\right)$ , where the  $a_n$  ( $n = 1, 2, 3, \dots$ ) are constants.

(i) Verify that the series in (d) satisfies the PDE and conditions (a) and (c).

(8 marks)

(ii) Find the  $a_n$  so that (b) is satisfied.

(9 marks)

(iii) Deduce that the potential energy stored in the string at time  $t$  is

$$\frac{16\rho h^2 c^2}{\pi^4 l} \sum_{m=0}^{\infty} \frac{\cos^2 \left\{ \frac{(2m+1)\pi ct}{l} \right\}}{(2m+1)^4}.$$

(8 marks)

- 3 In a compressible, static and uniform gas, with constant density  $\rho_0$  and pressure  $p_0$ , due to the passage of a sound disturbance, there are *small* changes in density  $\rho$  and pressure  $p$ .

- (i) Given that the exact equation of continuity is

$$\rho_t + (\rho u)_x + (\rho v)_y + (\rho w)_z = 0,$$

where  $\mathbf{u}(\mathbf{x}, t) = u(\mathbf{x}, t)\mathbf{i} + v(\mathbf{x}, t)\mathbf{j} + w(\mathbf{x}, t)\mathbf{k}$  is the velocity field of the perturbed state, obtain a valid approximation to the exact continuity equation in the limit of linear theory (i.e. small changes).

(5 marks)

- (ii) Using Newton's Second Law, again in linear approximation, and given that  $p$  is a function of  $\rho$ , show that

$$\rho_{tt} = c^2(\rho_{xx} + \rho_{yy} + \rho_{zz}),$$

where  $c^2$  is a constant which should be defined.

(10 marks)

- (iii) In a particular case  $\rho = \rho_0(1 + s)$  and supposing a potential flow (i.e.  $u = \phi_x$ ,  $v = \phi_y$  and  $w = \phi_z$ , where  $\phi$  is the velocity potential) show that for the linearised continuity equation

$$s_t = -(\phi_{xx} + \phi_{yy} + \phi_{zz}).$$

Provided that all disturbances decay to a steady state as  $|\mathbf{x}| \rightarrow \infty$ , deduce that

$$c^2 s = -\phi_t,$$

and hence show that

$$\phi_{tt} = c^2(\phi_{xx} + \phi_{yy} + \phi_{zz}),$$

i.e. the velocity potential also satisfies the three-dimensional form of the wave equation.

(10 marks)

- 4 The equilibrium position of the free surface of a liquid of infinite depth is  $z = 0$ , where  $z$  is measured vertically upwards. A surface wave causes the displacement of this surface to be  $\eta(x, t)$ , where  $x$  is measured along the undisturbed surface and

$$\eta = a \cos(kx - \omega t),$$

with  $a$ ,  $k$  and  $\omega$  being positive constants with  $a$  small. You are given that the velocity potential  $\phi = \phi(x, z, t)$  satisfies

$$\phi_{xx} + \phi_{zz} = 0.$$

You are also given that (a)  $\phi_z \rightarrow 0$  as  $z \rightarrow -\infty$ ; (b)  $\phi_z = \eta_t$  at  $z = 0$ ; (c)  $\phi_t + g\eta = 0$  at  $z = 0$ .

- (i) Give a brief physical interpretation of (a), (b) and (c).

**(6 marks)**

- (ii) Find  $\phi(x, z, t)$  and show that  $\omega^2 = gk$ .

**(13 marks)**

- (iii) Determine the phase velocity  $c$  and the group velocity  $c_g$  in terms of  $k$ . Show that  $c_g = c/2$ , and state what is the speed of the propagation of energy.

**(6 marks)**

- 5 (i) Solve, using the method of characteristics, the equation

$$yz_x + xz_y = xy,$$

given that  $z = e^{-y^2}$  on  $x = 0$  for  $y \geq 0$  and that  $z = e^{-x^2}$  on  $y = 0$  for  $x \geq 0$ . [We use the notation  $z_x = \frac{\partial z}{\partial x}$  etc.]

**(13 marks)**

- (ii) On what region  $D$  in the  $x$ - $y$  plane is the solution defined uniquely? Verify that  $z$  is everywhere continuous in  $D$ , but that  $z_x$  and  $z_y$  are discontinuous across one curve. Determine the curve of discontinuity.

**(7 marks)**

- (iii) More generally, suppose that within a region  $D^*$  in the  $x$ - $y$  plane, the solution  $z = z(x, y)$  of

$$Pz_x + Qz_y = R,$$

where  $P$ ,  $Q$ ,  $R$  are continuous functions of  $x$ ,  $y$ ,  $z$ , is everywhere continuous, but that there may be discontinuities in  $z_x$  and  $z_y$  across a curve  $\Gamma$ . Show that  $dy/dx = Q/P$  on  $\Gamma$ .

**(5 marks)**

**End of Question Paper**